



look good. feel good.
get into shape.

About ACSM's Current Comments

Current Comments are official statements by the American College of Sports Medicine concerning topics of interest to the public at large.

Permission to reprint text content

Permission to reprint this American College of Sports Medicine "Current Comment" contingent upon the article being reprinted in-total and without alteration, and with the printing of the following citation on each page or Web screen: "Reprinted with permission of the American College of Sports Medicine, "Alcohol and Athletic Performance," April 2000 www.acsm.org.

Cardiovascular system

Aerobic/endurance training can help to maintain and improve various aspects of heart and lung function and cardiac output, and such exercise can enhance endurance. Strength/resistance training will help offset the loss in muscle mass and strength typically associated with aging, thereby improving functional capacity. Also important, reduction in risk factors associated with disease (heart disease, diabetes, osteoporosis, and so on) will improve health status and contribute to an increase in lifespan. Together, these training adaptations will greatly improve the functional capacity of older men and women, therefore improving their quality of life and extend independent living.

Effects of Aerobic Exercise Training: Over the past ten years, we have learned that older persons can adapt to a program of regular aerobic training as well as their younger counterparts. Older adults can achieve the same 10 to 30 percent increase in VO₂max in response to endurance exercise training as young adults. The magnitude of these adaptations in VO₂max in older adults is a function of training intensity; low intensity training elicits only marginal changes. The increase in VO₂max in older adults is a result of improvements in both maximal cardiac output and a-v O₂ difference. In addition, improvements in submaximal endurance capacity and the greater ability to tolerate higher levels of physical activity are important training adaptations. For improvements in cardiovascular fitness, the American College of Sports Medicine recommends an exercise intensity of 55/65 to 90 percent of maximum heart rate (or 40/50 to 80 percent of heart rate reserve). ACSM further recommends accumulating 20-60 minutes at that level three to five days a week. The lower ranges are for unfit or even frail individuals who are about to begin an exercise program.

ACSM... Advancing Health.

**Through Science
Education, and
Medicine.**



**AMERICAN COLLEGE
of SPORTS MEDICINE
www.acsm.org**

401 W. Michigan St., Indianapolis, IN 46202-3233 USA
317.637.9200 tel 317.634.7817 fax

www.acsm.org/current_comments



**AMERICAN COLLEGE
of SPORTS MEDICINE
www.acsm.org**

Current Comments

*Report on Exercise and
the Older Adult*



prescription for prevention

Current evidence clearly indicates that participation in a regular exercise program is an effective way to reduce and/or prevent a number of the functional declines associated with aging. Older adults have the ability to adapt and respond to both endurance and strength training.

Aerobic/endurance training can help to maintain and improve various aspects of heart and lung function and cardiac output, and such exercise can enhance endurance. Strength/resistance training will help offset the loss in muscle mass and strength typically associated with aging, thereby improving functional capacity. Also important, reduction in risk factors associated with disease (heart disease, diabetes, osteoporosis, and so on) will improve health status and contribute to an increase in lifespan. Together, these training adaptations will greatly improve the functional capacity of older men and women, therefore improving their quality of life and extend independent living.

Written for ACSM by Robert S. Mazzeo, Ph.D., FACSM

ACSM Current Comment

Current evidence clearly indicates that participation in a regular exercise program is an effective way to reduce and/or prevent a number of the functional declines associated with aging. Older adults have the ability to adapt and respond to both endurance and strength training.

Decreased bone density is more common among older adults, evidence suggests that participation in regular exercise improves bone health and thus reduces the risk for developing osteoporosis. Further, this can reduce the incidence of breaks and fractures associated with falls.



Health Benefits

Most risk factors associated with disease increase with age, so the benefits of regular exercise are significant from a health perspective. Health benefits associated with cardiovascular disease risk factors include favorable changes in lipid profile, blood pressure, and body composition. Older adults can improve their plasma lipoprotein lipid profiles with exercise training similar to those observed in younger adults and may include modest increases in plasma HDL levels and reductions in LDL and plasma triglyceride levels. This results in more favorable HDL/LDL and light cholesterol: HDL ratios.

Exercise and the Older Adult

An ACSM Report

By the year 2030, the number of individuals 65 years and over will reach 70 million in the United States alone. Those 85 and older will then be the fastest-growing segment of our population. We must determine the extent and mechanisms by which exercise and physical activity can improve health, functional capacity, quality of life, and independence in this population.

Aging and Exercise

Aging is a multi-faceted process in which a variety of factors interact (genetics, lifestyle, disease) and frequently is associated with reduced functional capacity and chronic illness. In addition, physical inactivity and maintenance of a sedentary lifestyle represent a significant health risk to aging individuals. Older adults can safely participate in regular exercise programs (aerobic and strength). Regular physical activity has been shown to elicit a number of favorable adaptations that contribute to healthy aging. Further, the trainability of older individuals is evident from their ability to adapt and respond to both endurance and strength training.

Endurance training can help to maintain and improve various aspects of cardiovascular function as measured by VO2 max, cardiac output and a VO2 difference as well as enhance submaximal performance. It is important to note that reductions in risk factors associated with disease states (heart disease, diabetes, etc.) will improve health status and contribute to an increase in lifespan. Strength training will help offset the loss in muscle mass and strength typically associated with normal aging.

Additional benefits include improved bone health and thus reductions in risk for osteoporosis; improved postural stability, reducing the risk of falling; and increased flexibility and range of motion. Together, these benefits associated with regular exercise and physical activity will contribute to a healthier, more independent lifestyle, greatly improving functional capacity and quality of life for the fastest-growing segment of our population.



effects of resistance training

Given an adequate training stimulus, older adults can make significant gains in strength. A two- to threefold increase in strength can be accomplished in three to four months in fibers recruited during training in older adults. With more prolonged resistance training, even a modest increase in muscle size is possible. Because sarcopenia and muscle weakness are so prevalent in the aging population, it is important to devise strategies for preserving or increasing muscle mass in the older adult. With increasing muscle strength come increased levels of spontaneous activity in both healthy, independent older adults and very old and frail men and women. Strength training, in addition to its possible effects on insulin action, bone density, energy metabolism, and functional status, is also an important way to increase levels of physical activity in the older adult.

Effects of Aging

Loss of muscle mass (sarcopenia) with age in humans is well documented. A primary factor in sarcopenia is disuse of skeletal muscle, resulting in atrophy. A reduction in muscle strength is directly associated with loss of muscle mass. Inactivity may also play a role, contributing to other factors affecting aging muscle mass, including:

- Neuromuscular realignment (changes in motor units and innervation of fibers)
- Reduction in growth factors
- Changes in muscle protein turnover
- Post-exercise: Rehydrate first and consume food to retard any alcohol absorption

The consequences of sarcopenia can be extensive; individuals are more susceptible to falls and fractures, impaired in ability to regulate body temperature, slower in metabolism, possibly deficient in glucose regulation and may suffer an overall loss in the ability to perform everyday tasks. Muscle atrophy appears to result from a gradual loss of both muscle fiber size and number. A gradual loss in muscle cross-sectional area is consistently found with advancing age; by age 50, about ten percent of muscle area is gone. After 50 years of age, the rate of accelerates significantly. Muscle strength declines by approximately 15 percent per decade in the sixties and seventies and by about 30 percent thereafter. Although intrinsic muscle function is reduced with advancing age, age-related decrease in muscle mass is responsible for almost all loss of strength in the older adult. The number of functional motor units also declines with advancing age, which requires surviving motor units to innervate a greater number of muscle fibers.

Positive Effects of Exercise

Aerobic training reduces resting blood pressure in hypertensive young persons, and this training effect is evident in older hypertensive adults as well. Changes in body composition are associated with aerobic exercise training in older populations. A modest yet significant reduction in total percent body fat is generally observed with exercise training in older adults, which can occur despite a maintained body weight. It is important to note that in older men, a decrease in intra-abdominal fat accounts for the greatest relative loss of fat mass. This finding is significant in that intra-abdominal fat increases the most with advancing age and is associated with cardiovascular disease, particularly in men. In addition to reducing heart disease risk, regular exercise results in an increase in insulin sensitivity in older adults. As insulin resistance increases with age, the positive effects of regular aerobic exercise in older individuals on improving insulin sensitivity and increasing glucose transporters in muscle are of clinical importance for the treatment and prevention of adult-onset diabetes.

